

EFI 93-43
TECHNION-PH-93-31
hep-ph/9308308
August 1993

Framework for Identification of Neutral B Mesons

Michael Gronau

*Department of Physics, Technion-Israel Institute of Technology
Technion City, 32000 Haifa, Israel*

and

Jonathan L. Rosner

*Enrico Fermi Institute and Department of Physics
University of Chicago, Chicago, Illinois 60637*

ABSTRACT

We introduce a method for the study of CP-violating asymmetries in tagged states of neutral B mesons with arbitrary coherence properties. A set of time-dependent measurements is identified which completely specifies the density matrix of the initial state in a two-component space with basis vectors B^0 and $\bar{B}^0$, and permits a determination of phases in the Cabibbo-Kobayashi-Maskawa matrix. For a given tagging configuration, the measurement of decays both to flavor eigenstates and to CP eigenstates provides the necessary information.

The search for CP violation in systems involving B mesons is being pursued using several approaches. Decay modes for which a CP-violating rate asymmetry would be particularly easy to interpret include those of neutral B mesons to CP eigenstates such as $J/\psi K_S$ and $\pi^+\pi^-$. However, one is required to identify the flavor of the decaying meson: was it a $B^0(=\bar{b}d)$ or $\bar{B}^0(=b\bar{d})$ at the time of production?

One suggestion for neutral B meson identification involves the reaction

$$e^+ + e^- \rightarrow \Upsilon(4S) \rightarrow B^0 + \bar{B}^0 \quad , \quad (1)$$

in which the flavor of one meson is identified by correlating it with the semileptonic decay products of the other. In this configuration, with $C(B\bar{B}) = -1$, any CP-violating asymmetry is odd in the difference of times of the two decays [1,2], so asymmetric storage rings may be needed to provide the necessary time resolution. It has also been proposed [2] to produce the $B\bar{B}$ pair at a slightly higher energy, in association with an extra photon from B^* decay, in order to produce a state with $C(B\bar{B}) = 1$ to overcome this difficulty. In both cases the leptonic “tag” of a neutral B gives rise to an associated pure B^0 or $\bar{B}^0$.

Another method for determining the flavor of a neutral B relies on the semileptonic decay of a meson or baryon containing a b quark produced at high energies in association with the neutral B and with many other particles. Here the flavor of the decaying B is misidentified part of the time as a result of $B^0 - \bar{B}^0$ mixing, but in contrast to the case of production at and just above the $\Upsilon(4S)$ resonance, the initial state of the decaying neutral B is usually assumed to be an *incoherent* mixture of particle and antiparticle [3].

In order to avoid having to measure the decay of the associated particle, it has been proposed [4] to study the correlation of the decaying neutral B with charged pions produced nearby in phase space. This method has been used to identify neutral D mesons through the decays $D^{*+} \rightarrow \pi^+ D^0$ or $D^{*-} \rightarrow \pi^- \bar{D}^0$ [5]. As in semileptonic tagging at high energy, the initial B^0 and $\bar{B}^0$ states were assumed in Ref. [4] to be incoherent with one another.

In the present paper we develop a way to identify a neutral B meson which embraces the above examples but allows one to test experimentally for arbitrary coherence properties of the initial state. The method relies upon the description of the initial tagged state in terms of a density matrix, in a two-component “quasispin” space spanned by B^0 and $\bar{B}^0$. The two-component description is standard for kaons (see, e.g., Refs. [6]), and has been applied as well to the case of B ’s [7].

We have found a set of time-dependent measurements of decays of a tagged B which completely specifies the density matrix, permitting one to obtain phases in the Cabibbo-Kobayashi-Maskawa (CKM) matrix through a study of CP-violating decay rate asymmetries. Each tagging procedure specifies a different density matrix. Previously it was necessary to anticipate the properties of such a matrix

with the help of theoretical assumptions. Our result avoids the need for any such procedure, and provides an answer to the question of coherence of the initial state.

In brief, we advocate the measurement of B decays both to states of identifiable flavor (such as $J/\psi K^{*0}$, with $K^{*0} \rightarrow K^+ \pi^-$) and to eigenstates of CP (such as $J/\psi K_S$). The observables which specify the initial density matrix are: (1) a modulation amplitude of the time-dependent term; (2) a shift in phase of the time-dependent term, and (3) an overall intensity associated with the production of the CP eigenstate. Once these quantities are measured, one not only determines the density matrix, but also obtains valuable information about weak phases.

We now define the basis states and the density matrix. We choose

$$|B^0\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad , \quad |\overline{B}^0\rangle = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \quad . \quad (2)$$

A density matrix ρ allows one to discuss incoherent and coherent states in a unified manner. The most general such 2×2 matrix has the form

$$\rho = \frac{1}{2} [1 + \mathbf{Q} \cdot \boldsymbol{\sigma}] \quad , \quad (3)$$

where $\mathbf{Q}$ is a vector describing polarization in quasispin space, satisfying $\mathbf{Q}^2 \leq 1$, and σ_i ($i = 1, 2, 3$) are the Pauli matrices. We give two examples:

(1) Any pure state corresponds to a linear combination of B^0 and $\overline{B}^0$ with arbitrary complex coefficients, whose sum of absolute squares equals unity. Such a state can be denoted by a density matrix with $Q \equiv |\mathbf{Q}| = 1$.

(2) An arbitrary incoherent combination of B^0 and $\overline{B}^0$ with relative probabilities P_1 and $P_2 = 1 - P_1$ corresponds to a diagonal density matrix with $Q_1 = Q_2 = 0$, $Q_3 = 2P_1 - 1$. This is the case we considered to hold in Ref. [4].

As a special case of either (1) or (2), one describes the density matrices for initial B^0 and $\overline{B}^0$ by $\text{diag}(1,0)$ and $\text{diag}(0,1)$, respectively.

The probability for a transition from an initial state denoted by the density matrix ρ_i to a final state denoted by ρ_f is then

$$I(f) = \text{Tr} (\rho_i T^\dagger \rho_f T) \quad , \quad (4)$$

where T is the amplitude which time-evolves the state from i to f . Here we shall take f to denote an arbitrary coherent superposition of B^0 and $\overline{B}^0$ at time t . We shall also be able to write a density matrix ρ_f which takes account of the *decay* of this superposition.

The time-evolution can be visualized by transforming from the $B^0, \overline{B}^0$ basis to that of mass eigenstates. We shall neglect differences between lifetimes of these two eigenstates [1], and denote them by B_L (“light”) and B_H (“heavy”):

$$|B_L\rangle = p|B^0\rangle + q|\overline{B}^0\rangle \quad , \quad |B_H\rangle = p|B^0\rangle - q|\overline{B}^0\rangle \quad . \quad (5)$$

Here p and q are approximately of unit magnitude [1]. We adopt a phase convention in which the $B^0 - \bar{B}^0$ mixing amplitude is real, so that $p = q = 1/\sqrt{2}$. This differs from a more standard convention [1] by a phase which we shall take into account when calculating amplitudes for decays of b quarks. The transformation between flavor eigenstates and mass eigenstates is then implemented by the unitary matrix

$$U \equiv \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \frac{1}{\sqrt{2}}(\sigma_1 + \sigma_3) \quad . \quad (6)$$

The matrix describing the time evolution is diagonal in the mass eigenstate basis:

$$e^{-iM_D t} \equiv e^{-\Gamma t/2} \text{diag}(e^{-im_L t}, e^{-im_H t}) = e^{-\Gamma t/2} e^{-i\bar{m}t} e^{i\sigma_3 \Delta m t/2} \quad , \quad (7)$$

where $\bar{m} \equiv (m_H + m_L)/2$, $\Delta m \equiv m_H - m_L$. Thus the time evolution operator T in the $B^0, \bar{B}^0$ basis is just $T = U^\dagger e^{-iM_D t} U = e^{-\Gamma t/2} e^{-i\bar{m}t} e^{i\sigma_1 \Delta m t/2}$.

The trace for the transition probability $I(f)$ can be computed by applying the matrices U and $U^\dagger$ to the initial and final density matrices instead of to $e^{-iM_D t}$. Defining $\rho' \equiv U \rho U^\dagger$, we find that the effect of U is to rotate $\mathbf{Q}$ into $\mathbf{Q}'$, where

$$Q'_1 = Q_3 \quad , \quad Q'_2 = -Q_2 \quad , \quad Q'_3 = Q_1 \quad . \quad (8)$$

We already remarked that states which are pure B^0 or pure $\bar{B}^0$ correspond to $Q_3 = \pm 1$, $Q_1 = Q_2 = 0$. Their transformed density matrices are $\rho'_f = (1/2)(1 \pm \sigma_1)$, respectively, since then $Q'_1 = \pm 1$, $Q'_2 = Q'_3 = 0$. The transition probability can now be written in terms of traces as

$$I(f) = \text{Tr} (\rho'_i e^{iM_D^* t} \rho'_f e^{-iM_D t}) \quad . \quad (9)$$

The states B^0 and $\bar{B}^0$ at the time of decay t may be identified by their decays to states of identifiable flavor, e.g., $B^0 \rightarrow J/\psi K^{*0}$, with $K^{*0} \rightarrow K^+ \pi^-$. We assume that a single weak subprocess contributes to the decay, which is an excellent approximation for these final states [1].

With the convention in which the mixing amplitudes p and q in the neutral B mass eigenstates are real, the weak decay amplitudes for $B^0 \rightarrow J/\psi K^{*0}$ and $\bar{B}^0 \rightarrow J/\psi \bar{K}^{*0}$ may be denoted $Ae^{-i\beta}$ and $Ae^{i\beta}$, respectively, where $\beta = \text{Arg}(-V_{cb}^* V_{cd}/V_{tb}^* V_{td})$, and V_{ij} are elements of the Cabibbo-Kobayashi-Maskawa matrix specifying the charge-changing weak couplings of quarks. We find

$$I \left(\begin{array}{c} B^0 \\ \bar{B}^0 \end{array} \right) = \frac{1}{2} |A|^2 e^{-\Gamma t} [1 \pm (Q'_1 \cos \Delta m t - Q'_2 \sin \Delta m t)] \quad . \quad (10)$$

No reference to Q'_3 appears. The sine and cosine terms may be combined into a cosine of a phase-shifted argument by defining

$$Q'_1 \equiv Q'_\perp \cos \delta \quad , \quad Q'_2 \equiv Q'_\perp \sin \delta \quad , \quad (11)$$

so that

$$I \left(\frac{B^0}{\bar{B}^0} \right) = \frac{1}{2} |A|^2 e^{-\Gamma t} [1 \pm Q'_\perp \cos(\Delta m t + \delta)] \quad . \quad (12)$$

Whenever the initial state is any incoherent mixture of B^0 and $\bar{B}^0$ with relative probabilities P_1 and $P_2 = 1 - P_1$, respectively, one simply sets $Q'_\perp = 2P_1 - 1$ and $\delta = 0$ in the above expression.

We consider a charge-symmetric production process (such as $\bar{p}p$ or e^+e^- collisions) in which an arbitrary coherent, partially coherent, or incoherent combination of neutral B^0 and $\bar{B}^0$ is produced, with an additional particle of specific charge (such as a charged lepton or pion) bearing some specific kinematic relation to it. This relation could consist, for example, of a lepton with specific transverse momentum or range of transverse momenta, a definite-mass combination $M(B\pi)$ (or range of such combinations), a specific angle or range of angles between the B and tagging particle, or a specific rapidity difference (or range thereof) between the two particles. It is necessary to determine the relation of this process to the one in which the charge of the tagging particle is the opposite.

Under the phase convention we have chosen for B^0 and $\bar{B}^0$, if we take $CP|B_L\rangle = |B_L\rangle$ and $CP|B_H\rangle = -|B_H\rangle$, the charge conjugation operation has the phase

$$C|B^0\rangle = -|\bar{B}^0\rangle \quad ; \quad C|\bar{B}^0\rangle = -|B^0\rangle \quad . \quad (13)$$

Under charge conjugation, the first and second rows and columns of the density matrix ρ are interchanged, so that $Q_1 \rightarrow Q_1$, $Q_2 \rightarrow -Q_2$, $Q_3 \rightarrow -Q_3$. We then find that the effect of charge conjugation is a rotation by π about the $3'$ direction:

$$C : \quad Q'_1 \rightarrow -Q'_1, \quad Q'_2 \rightarrow -Q'_2, \quad Q'_3 \rightarrow Q'_3 \quad . \quad (14)$$

Therefore, the decay rates $\bar{I}(f)$ for states tagged with antiparticles are then given in terms of those $I(f)$ for states tagged with particles by

$$\bar{I}(f; Q'_1, Q'_2, Q'_3) = I(f; -Q'_1, -Q'_2, Q'_3) \quad . \quad (15)$$

For a final state identified as a B^0 by its decay to $J/\psi K^{*0}$, the time-dependent asymmetry is

$$A(J/\psi K^{*0}) \equiv \frac{I(J/\psi K^{*0}) - \bar{I}(J/\psi K^{*0})}{I(J/\psi K^{*0}) + \bar{I}(J/\psi K^{*0})} = Q'_\perp \cos(\Delta m t + \delta) \quad . \quad (16)$$

As a consequence of the assumed charge symmetry of the production process, one has $\bar{I}(J/\psi K^{*0}) = I(J/\psi \bar{K}^{*0})$ and $\bar{I}(J/\psi \bar{K}^{*0}) = I(J/\psi K^{*0})$.

We can then measure both components Q'_1 and Q'_2 , or, equivalently, $Q'_\perp$ and δ , using decays to flavor eigenstates. The ALEPH Collaboration [8] has recently measured a time-dependent asymmetry of the above form, fitting it under the

assumption $\delta = 0$. It should be possible to set limits on δ from this measurement. A way of measuring Q'_3 will be described when we come to discuss CP-violating asymmetries.

The above expressions are quite general and not restricted to any specific tagging method. The initial density matrix ρ can refer to any method of preparation, including specific configurations at or just above the $\Upsilon(4S)$ resonance in e^+e^- collisions.

A measurement of Q'_3 for neutral nonstrange B mesons can be performed by utilizing their decays to the specific CP eigenstate $J/\psi K_S$. In our phase convention, the amplitudes for B^0 and $\bar{B}^0$ to decay into $J/\psi K_S$ are $A'e^{-i\beta}/\sqrt{2}$ and $-A'e^{i\beta}/\sqrt{2}$. Here we have taken into account the intrinsic negative CP of the $J/\psi K_S$ state, and neglected the small CP violation in the kaon system. The density matrix for the final state is then

$$\rho_{J/\psi K_S} = \frac{1}{2}|A'|^2 \begin{bmatrix} e^{-i\beta} & -e^{i\beta} \\ -e^{-i\beta} & \end{bmatrix} = \frac{1}{2}|A'|^2 \begin{bmatrix} 1 & -e^{2i\beta} \\ -e^{-2i\beta} & 1 \end{bmatrix} \quad . \quad (17)$$

For reference, we also note that

$$\rho_{J/\psi K_L} = \frac{1}{2}|A'|^2 \begin{bmatrix} e^{-i\beta} & e^{i\beta} \\ e^{-i\beta} & \end{bmatrix} = \frac{1}{2}|A'|^2 \begin{bmatrix} 1 & e^{2i\beta} \\ e^{-2i\beta} & 1 \end{bmatrix} \quad . \quad (18)$$

The expressions for the decay rates for states prepared with particle and antiparticle tags are then

$$\begin{aligned} I\left(J/\psi \begin{array}{c} K_L \\ K_S \end{array}\right) &= \frac{1}{2}|A'|^2 e^{-\Gamma t} \{1 \pm [Q'_3 \cos 2\beta + Q'_\perp \sin 2\beta \sin(\Delta mt + \delta)]\}, \\ \bar{I}\left(J/\psi \begin{array}{c} K_L \\ K_S \end{array}\right) &= \frac{1}{2}|A'|^2 e^{-\Gamma t} \{1 \pm [Q'_3 \cos 2\beta - Q'_\perp \sin 2\beta \sin(\Delta mt + \delta)]\}. \end{aligned} \quad (19)$$

As in the case of decays to the flavor eigenstates, the time-dependent term has a phase shift δ and a modulation amplitude $Q'_\perp$. It depends upon $\sin(\Delta mt + \delta)$ rather than $\cos(\Delta mt + \delta)$.

The decay asymmetry for the $J/\psi K_S$ final state is then

$$A(J/\psi K_S) \equiv \frac{I(J/\psi K_S) - \bar{I}(J/\psi K_S)}{I(J/\psi K_S) + \bar{I}(J/\psi K_S)} = \frac{-Q'_\perp \sin 2\beta \sin(\Delta mt + \delta)}{1 - Q'_3 \cos 2\beta} \quad . \quad (20)$$

The component Q'_3 (which appears even in the absence of CP violation) is a necessary ingredient in the discussion of possible coherence. It is this component that leads to correlations between K_S and K_L produced in ϕ decay, as discussed in Ref. [9]. In order to learn its value, we measure the rate for $J/\psi K_S$ production (summing over particle and antiparticle tags, so that the time-dependent terms

cancel). We compare this with the corresponding sum of rates (which also has no time-dependence, and is independent of Q'_3) for production of a flavor eigenstate K^0 .

Since we cannot observe a K^0 directly, we must resort to indirect means. If the rate of production of charged and neutral B 's is equal, as is expected to be the case for high-energy e^+e^- collisions [4], one can compare the rates for $J/\psi K^+$ and $J/\psi K_S$ production, making use of the fact that the decays of B mesons to $J/\psi K$ involve the quark subprocess $b \rightarrow c\bar{c}s$, which conserves isospin [10].

A more general method makes use of the expectation that the K^0/K^+ ratio will be the same as the K^{*0}/K^{*+} ratio. The flavor eigenstates involving K^{*0} , K^+ , and K^{*+} are all observable. Thus we can determine the ratio $|A'/A|^2$.

In either method we learn the relative normalization of rates for decays to flavor eigenstates and CP eigenstates. We thereby determine the magnitude of the term $Q'_3 \cos 2\beta$, and then use the asymmetry in $J/\psi K_S$ decay to measure $\sin 2\beta$. With the possibility of a discrete ambiguity (unlikely for known ranges of CKM parameters), we then obtain $\cos 2\beta$, thereby finding Q'_3 itself.

The corresponding decay asymmetry for the $\pi^+\pi^-$ final state is easily calculated by our method. We neglect penguin effects [11], which can be dealt with by studying the $2\pi^0$ final state. With our phase convention for b quarks, the result can be obtained by the substitution $\beta \rightarrow -\alpha$ in the corresponding result for the $J/\psi K_L$ final state, where $\alpha = \text{Arg}(-V_{tb}^* V_{td}/V_{ub}^* V_{ud})$. We find

$$A(\pi^+\pi^-) \equiv \frac{I(\pi^+\pi^-) - \bar{I}(\pi^+\pi^-)}{I(\pi^+\pi^-) + \bar{I}(\pi^+\pi^-)} = \frac{-Q'_\perp \sin 2\alpha \sin(\Delta mt + \delta)}{1 + Q'_3 \cos 2\alpha} \quad . \quad (21)$$

Since we have already measured all components of $\mathbf{Q}'$ and the phase δ , this result can be used to extract α .

To sum up, we have developed a density-matrix formalism for the discussion of decays of neutral B mesons which have been “tagged” in any arbitrary fashion. The density matrix is specified in terms of a polarization vector $\mathbf{Q}$ in the basis labelled by B^0 and $\bar{B}^0$, or $\mathbf{Q}'$ in the basis of mass eigenstates. It is the components of this polarization vector which determine all observable CP-violating asymmetries, and which can be learned by measurements on the time-dependence of decays to $J/\psi K^*$ and $J/\psi K$. One learns as well the angle β . Decays to final states such as $\pi^+\pi^-$ can then provide information on other phases of the Cabibbo-Kobayashi-Maskawa matrix.

We are grateful to B. Kayser, H. Lipkin, A. I. Sanda, and S. Stone for helpful discussions, and to D. G. Cassel and H. Tye for extending the hospitality of the Newman Laboratory of Nuclear Science, Cornell University, during part of this investigation. This work was supported in part by the United States – Israel Binational Science Foundation under Research Grant Agreement 90-00483/2, by the Fund for Promotion of Research at the Technion, and by the U. S. Department of Energy under Grant No. DE FG02 90ER-40560.

REFERENCES

- [1] I. Dunietz and J. L. Rosner, Phys. Rev. D**34**, 1404 (1986); I. Dunietz, Ann. Phys. (N.Y.) **184**, 350 (1988); I. Bigi and A. I. Sanda, Nucl. Phys. **B281**, 41 (1987); Y. Nir and H. Quinn, Ann. Rev. Nucl. Part. Sci. **42**, 211 (1992).
- [2] *Updated Proposal for a B Factory Upgrade for the Cornell Electron Storage Ring*, Cornell University report CNLS-CESR-B-UPD-PROP, May, 1993 (unpublished); *Progress Report on Physics and Detector at KEK Asymmetric B Factory*, KEK Report 92-3, May 1992 (unpublished); *An Asymmetric B Factory Based on PEP*, Lawrence Berkeley Laboratory LBL PUB-5303, February, 1991 (unpublished).
- [3] See, e.g., *B Physics at Hadron Accelerators*, Proceedings of a workshop at Fermilab, Nov. 16–17, 1992, edited by J. A. Appel (Fermilab, Batavia, IL, 1992); *Beauty 93* (Proceedings of the First International Workshop on B Physics at Hadron Machines, Liblice Castle, Melnik, Czech Republic, Jan. 18–22, 1993), edited by P. E. Schlein, Nucl. Instrum. Meth. **331**, No. 1 (1993); M. Chaichian and A. Fridman, Phys. Lett. **B298**, 218 (1993).
- [4] M. Gronau, A. Nippe, and J. L. Rosner, Phys. Rev. D**47**, 1988 (1993).
- [5] S. Stone, in *Heavy Flavors*, edited by A. J. Buras and M. Lindner (World Scientific, Singapore, 1992).
- [6] J. S. Bell and J. Steinberger, in *Proceedings of the Oxford International Conference on Elementary Particles*, 19–25 September, 1965, edited by T. R. Walsh *et al.* (Rutherford High Energy Laboratory, Chilton, England, 1966), p. 195; T. D. Lee and C. S. Wu, Ann. Rev. Nucl. Sci. **16**, 511 (1966).
- [7] H. J. Lipkin, Phys. Lett. **B219**, 474 (1989); Weizmann Institute preprint WIS-93-12, 1993 (unpublished).
- [8] ALEPH Collaboration, CERN report CERN-PPE/93-99, June, 1993, submitted to Phys. Lett. B.
- [9] H. J. Lipkin, Phys. Rev. **176**, 1715 (1968).
- [10] H. J. Lipkin and A. I. Sanda, Phys. Lett. **B201**, 541 (1988).
- [11] M. Gronau, Phys. Rev. Lett. **63**, 1459 (1989); Phys. Lett. **B300**, 163 (1993); M. Gronau and D. London, Phys. Rev. Lett. **65**, 3381 (1990).